

SPECIAL ARTICLE

# COVID-19 and pain: what we know so far

## COVID-19 y dolor: qué conocemos hasta ahora

### ABSTRACT:

SARS-CoV-2 is a new coronavirus spread throughout the world that has forced the declaration of a pandemic by the World Health Organization. Its symptomatology, which varies from flu symptoms to death, has resulted in a fracture in the life of the population, also affecting health care. We have carried out a systematic bibliographic search in the main online databases (PubMed, Medline, UpToDate, Embase, Scopus and Cochrane Library) with the following keywords: “COVID-19”, “coronavirus and SARS-CoV-2”, “pain” and “chronic pain”. Publications have been included from the description of the disease until December 2020, about the impact it has had on pain in its different areas, clinical characteristics and management.

### AUTHORS:

**María Teresa Fernández Martín**, Facultativo Especialista de Área. Servicio de Anestesia, Reanimación y Terapéutica del Dolor. Hospital Universitario Río Hortega. Valladolid

**Esperanza Ortigosa Solórzano**, Coordinadora de la Unidad del Dolor. Servicio de Anestesia, Reanimación y Terapéutica del Dolor. Hospital Universitario de Getafe. Madrid

### CORRESPONDENCE:

***eosolorzano@gmail.com***

### RESUMEN:

El SARS-CoV-2 es un nuevo coronavirus extendido por todo el mundo que ha obligado a la declaración de pandemia por la Organización Mundial de la Salud. Su sintomatología, que varía desde síntomas gripales hasta la muerte, ha supuesto una fractura en la vida de la población, afectando también a la atención sanitaria. Hemos realizado una búsqueda bibliográfica sistemática en las principales bases de datos *online* (PubMed, Medline, UpToDate, Embase, Scopus y Cochrane Library) con las siguientes palabras clave: “COVID-19”, “*coronavirus and SARS-CoV-2*”, “*pain*”, “*chronic pain*”. Se han incluido publicaciones desde la descripción de la enfermedad hasta diciembre de 2020, acerca de la repercusión que ha tenido sobre el dolor en sus diferentes ámbitos, las características clínicas y su manejo.

RECEIVED: 4 / January / 2021

ACCEPTED: 10 / January / 2021

**Key words:** COVID-19, SARS-CoV-2, pain, chronic pain, COVID management, COVID treatment.

**Palabras clave:** COVID-19, SARS-CoV-2, dolor, dolor crónico, manejo del COVID, tratamiento del COVID.

## Introduction

In late December 2019, a new disease of unknown etiology appeared in Wuhan (1), China. On January 30, 2020 the World Health Organization general director declared it as a Public Health Emergency of International Importance. The appearance of this new virus, SARS-CoV-2, known as COVID-19, caused a fracture in the way of living and working around the world, with effects in the future that are still unknown.

The outbreak in Wuhan (China), spread to Europe and America, imposed some strategies to contain the infection, such as social isolation, the absence of interpersonal relationships, cessation of commercial activity and non-essential services.

The high level of attention required by the number of patients affected by COVID-19 forced the redistribution of health resources, with the suspension of non-urgent activity. Although the right to health implies the right to adequate pain treatment (The Spanish Constitution of 1978, in its article 43, recognizes the right to health protection and therefore the right of citizens to receive health care adequate) and the World Health Organization (WHO) determined that pain relief is a fundamental right (2), under which circumstances it was considered a non-urgent procedure, so that the follow-up and treatment of chronic pain was decreased or interrupted during the pandemic. This fact has affected both treating patients and mental health.

The objective of this review was to analyze the impact of COVID-19 on chronic pain through bibliographic reviews. Systematic reviews and extension of meta-analyses have been selected for scope reviews (PRISMA-ScR). The databases consulted were PubMed, Scopus, UpToDate, Embase and the Cochrane Library. The unpublished literature search included: sites for pain organizations, the World Health Organization (WHO). The articles that met the inclusion criteria were those on the management of patients with chronic pain during the COVID-19 pandemic published after the pandemic (December 2019). All primary research studies, reviews, meta-analyses, guidelines and texts published on the sites were considered. There was no language restriction.

## COVID-19 as a factor involved in pain

In late December 2019, an outbreak of pneumonia cases of unknown etiology in the city of Wuhan in Hubei province, China, was announced and reported to the World Health Organization (WHO). It was quickly identified as a new betacoronavirus and linked to SARS-CoV and several other SARS-like coronaviruses transmitted by bats.

The initial symptoms to recognize the disease have been mainly fever, dry cough or dyspnea, although pain has also been an early symptom of this respiratory condition (sore throat, myalgia, low back pain, headache) (3,4). But this disease has shown a more dramatic side, some patients have suffered and suffer a torpid evolution, with ARDS (respiratory distress syndrome) that requires admission to the ICU and mechanical ventilation, resulting in death in 1.8-3.4 % of the cases (5).

Among all the symptoms we will focus our attention on pain. The IASP (International Association for the Study of Pain) defines pain, and especially chronic pain, as that persistent or recurrent pain for more than three months despite the resolution of tissue damage (6).

A good point to start this review is to remark that chronic pain, is the result of the interaction of biological, psychological and social factors, and the COVID-19 infection seems to involve each of them.

We know the relationship between acute viral disease (flu, SARS) with symptoms such as myalgia, fatigue, and general symptoms (7,8). After the SARS epidemic, patients with chronic fatigue syndrome, myalgia, depression and persistent sleep disorder up to 1-2 years after infection (9) have been observed.

Patients affected by other pathogens (Ros River Virus, Coxiella Burnetti and Epstein Barr virus) also experienced postviral fatigue syndromes, memory loss, and pain for more than a year (10).

Regarding SARS-CoV2, there are recent studies that specially focus on “the bradykinin storm” (11) and show a relationship between virus infection and inflammation. Bradykinin is not only involved in the pro-inflammatory state but also sensitizes the sensory fibers, which produces hyperalgesia (12). These inflammatory and immunological changes could explain why the elderly who are immunosuppressed and obese (increased inflammatory

*During the lockdown the elective pain consultations and invasive procedures were cancelled. This has led to an increase in pain and psychological worsening in these patients*

response) may have had a greater impact due to COVID-19. Additionally, COVID-19 infection may include encephalopathy, and structural changes in the brain and spinal cord like others viral infections (13).

The high number of cerebral complications in patients affected with SARS-CoV2, suggested that their neurotropism could cause neurodegenerative problems with an inflammatory basis (14). As shown above for other SARS-CoV infections, a mechanism of spread through the cribriform plate of the ethmoid bone, from the nose to the olfactory epithelium, has been proposed, where ACE2 receptors are highly expressed. This viral neurotropism through the olfactory bulb could be especially responsible for the frequently reported anosmia in these patients (15), SARS-CoV-2 can also reach deeper parts of the brain, including the thalamus and the brain stem, potentially contributing to respiratory failure (16) and to painful symptoms as well (Figure 1).

Brain PET metabolism was analyzed in comparison with healthy subjects throughout the brain, and the findings were confronted with the hypothesis of SARS-CoV-2 spread from the olfactory bulb to other limbic structures and possibly to the brain stem. Therefore, the hypometabolism of the radiopharmaceutical FDG could be considered a quantitative biomarker of this condition. However, cohort studies are needed to specify the relationship between hypometabolism and possible persistence of cognitive, emotional disorders, residual respiratory symptoms or painful complications (17).

In addition to the direct action of the virus, there are multiple factors that act or increase the painful symptoms. Chronic pain is associated with comorbidities and advanced age, which increases the risk of infections (including COVID-19). On the other hand, whether patients with chronic pain are more susceptible to infection is not yet entirely clear, although the interaction between pain and the immune system must be taken into account, since it has been shown that the persistence of pain and the absence of treatment induces immunosuppression (18).

Some treatments used to relieve pain can depress the immune system. Steroids that induce secondary adrenal failure (19). There are some studies that indicate that opioids decrease the natural killer (NK) cells, a dose-dependent effect, and interfere in the cellular response innate or indirectly by acting on the hypothalamic-pitui-




Figure 1.

tary-adrenal axis (producing corticosteroids) or in the sympathetic system (producing adrenaline). Both act on lymphocytes by negatively modulating the response of NK (20) cells. Morphine and fentanyl cause greater depression than buprenorphine (21). In animal models it has been shown that morphine can also act through the D1 pathway (dopamine receptors in the striated nucleus) (22).

Many patients who have been in the ICU have had long-term functional limitations, psychological disorders, and chronic pain. This suggests that patients who have been in the intensive care unit affected by this coronavirus will not behave differently (23).

This pandemic also has associated social aspects such as confinement, cessation of routine activities, financial impact, ... that have influenced the entire population. Isolation, limitation of interpersonal contact, fear of illness, uncertain future, economic ruin, etc., have increased the rates of fear and anxiety (24). Persistent stress can lead to varieties of severe mental illness (anxiety, depression, sleep disorder, or thoughts of suicide) (25).

It is important to recognize that thoughts and emotions are closely related to physical pain. Thus, the psychosocial conse-

quences of COVID-19 can influence physical pain and most likely vice versa as well. In other words, those patients who already had chronic pain will likely have a negative impact related to the infection, while those who did not previously have chronic pain can be added to this group (11) (Table I).

## Treatment during COVID-19: how we have changed

Patients with chronic pain require long-term multidisciplinary treatment even during a pandemic. They are mostly elderly with multiple comorbidities and are more susceptible to morbidity and mortality from COVID-19. It is necessary to review pain management practices during the COVID-19 era with respect to resource reallocation measures, community collaborations and the use of analgesics (steroids, opioids) and pain interventions. The patient with chronic pain faces a potential risk of functional and emotional deterioration during a pandemic, which can increase the long-term health burden (26).

During the pandemic lockdown, all elective pain consultations and interventional pain procedures were canceled or postponed (27). This interruption has had serious consequences, as it has led to an increase in pain, a psychological worsening and a decrease in the quality of life in these patients.

Compared to the general population, patients with chronic pain suffer from increased anxiety, stress and depression (28), which in turn increase the risk of chronic pain. Similarly, the relationship between sleep disturbance and pain is also bidirectional. Pain disturbs the quality of sleep and lack of sleep further exacerbates pain. Anxiety and sleep disturbance in confinement have increased uncertainty, worry and fear of illness, loneliness, and severe depressive symptoms. Aggravated by the fact that the isolation and closure of many services (rehabilitation, physiotherapy, psychological support groups etc.) have limited the mechanisms that many patients have been using to control pain.

Confinement has led the population, especially vulnerable patients with chronic pain, to reduced physical activity and exercise (caused by fear of leaving home and closure of sports centres), a change in diet (with increased consumption of “comfort” caloric foods and the absence of home help.

**Table I.**

1. Encephalopathy and structural changes in the brain and spinal cord could be included in Covid-19 infection
2. The neurotrophism of SARS-CoV-2 infection could cause neurodegenerative problems with an inflammatory base
3. Bradykinins contribute to pro-inflammatory state and also sensitize the sensitive fibers, leading to hyperalgesia
4. SARS-CoV-2 could contribute in the thalamus and brainstem to respiratory failure and painful symptoms

We can assure that all these changes have led to worsening comorbidities and chronic pain and that the interruption of treatment have had alarming consequences not only for individuals but for the health system and the whole society in both the short and long term.

In this scenario, telemedicine, which includes such diverse means of communication as telephone, video, email, fax, etc... has emerged as a key technology for efficient communication and a sustainable solution to provide essential health care services (Figure 2).


### Telemedicine a new tool

Chronic pain management requires frequent visits to the physician for both pharmacological and non-pharmacological treatment advice. Due to confinement, movement restrictions, social and physical distancing requirements, or the fear that health-care facilities may be infected, we have prevented patients from going to health-care facilities in person. This is where telemedicine has played a major role in helping patients to have a medical consult (29).

In accordance with clinical guidelines, recommendations, and practices, telemedicine should be considered whenever possible (30,31).

Telemedicine services offer many advantages, especially in routine or non-urgent care. Unnecessary and avoidable exposure of those involved in the provision of healthcare can be avoided. The need for personal protective equipment (PPE) is reduced leading to reduction in resource consumption. They improve access to health care and reduce media use across the already highly stressed health infrastructure during the current pandemic, all of which can provide substantial financial savings at all levels (32,33).

**Figure 2.**



Regarding the disadvantages currently there is a lack of clarity about the legal implications of its use. The use of technology cannot replace clinical medicine based on the in-person medical consultation and physical examination of the patient. Also, the possibilities of misdiagnosis and inability to carry out diagnostic radiological investigations are other limitations.

### What evidence do we have of its use in chronic pain?

The large-scale use of telemedicine in the treatment of chronic pain is rare and has generally focused on psychological interventions, exercise, stress reduction therapies etc. Several systematic reviews in adults report pain reduction, of disability, depression and anxiety in intervention groups compared to control groups (34,35). In contrast, psychological therapies performed via the Internet or mobile applications have shown only a small beneficial effect in children and adolescents with mixed chronic pain (36).

Chronic pain management is based on a good evaluation and medical history. This evaluation can be performed remotely using electronic versions of various questionnaires and scales. The reliability and validity of the telematic production of pain questionnaires such as the McGill questionnaire (in short format [SF-MPQ]) and the Pain Disability Index have been verified, with a significantly larger number of patients who prefer to conduct this type of tests electronically (37). Similarly, many questionnaires for the evaluation of patients with chronic lumbar pain have shown good reliability and moderate validity (38). The anxiety, depression and sleep questionnaires can also be collected electronically, so it is possible to perform a detailed pain assessment and history online.

The physical examination is the cornerstone of chronic pain management. A thorough physical examination is essential. Therefore, the first consultation should at least preferably be a face-to-face one, although in these special circumstances the patient must be made to understand that sometimes a complete physical examination cannot be performed. An important aspect of chronic pain management is non-drug treatment, such as ergonomics education, physical activity, lifestyle modification, counseling and coping skills, etc. And all these components can be easily delivered through telemedicine.

### Is telemedicine coming to stay with us?

One of the questions we should ask ourselves is whether telemedicine is a temporary tool in times of a pandemic or is it coming to be installed.

Although, according to the evidence, the beneficial effects of telemedicine are very promising, there remains substantial uncertainty surrounding many aspects of the studies conducted around it. These include small sample sizes, methodological defects, limited benefits, insufficient long-term evaluations, little collection of adverse events (39), etc. It is also necessary to explore the satisfaction and acceptance of patients, as well as studies of damage (produced in these) and desertions of this type of care.

If telemedicine is here to stay, which is highly likely, we must develop new clinical guidelines to apply in our medical practice. These guides should include general principles and a practical framework. The general principles should be common to all future guidelines, while the practical framework should mainly address the current scenario of the COVID-19 (40) pandemic (41,42). Among others, it should clearly specify the terms of the first consultation, follow-up consultations, patient management, consents, medications that can be prescribed through teleconsultation (43) etc. and the professionals who use this route must maintain the same ethical and professional standards as those observed in traditional face-to-face consultations.

### COVID inheritance

As far as we know so far there is not much data available on the impact of the COVID-19 pandemic and isolation measures on patients with chronic pain. What is clear is that the COVID-19 disease itself is associated with painful symptoms, including myalgia, arthralgia, abdominal pain, headache and chest pain, even in patients not admitted to the ICU (44). And survivors could have long-term effects, such as pain, that require our attention and management (45).

According to recent studies, more than 35 % of COVID-19 patients develop neurologic symptoms (46). All these neurological manifestations could lead to painful conditions, from peripheral neuralgia to post-traumatic pain syndrome, which must be diagnosed and treated, a challenge for doctors.

The study by Nieto et al. (47) on the impact that confinement has had on patients with chronic pain has shown changes in pain from the beginning of confinement, correlating pain intensity and disability scores and confirmed an increase in pain intensity, the frequency of episodes of pain, interference of pain (in daily activities, ability to work and leisure activities), distress caused by pain, and the effects of pain on sleep and physical activity. Specifically, when respondents were asked about changes (not necessarily related to pain), the mean scores showed a worsening in physical activity, social activities, work activities, emotional state, and overall well-being. These data highlight the importance of paying attention to people with chronic pain during health crises, as their pain problem and their overall health can worsen, as experts have pointed out.

Headache is the most common brain symptom, with a variable prevalence of 6.5 % to 23 % and an average prevalence of 8 % in different studies (48). Persistent daily cephalgia has been observed even weeks after recovery (49). Patients had a holocranial, oppressive cephalgia that began after 2 weeks after the recovery of respiratory symptoms, with no physical cause (venous thrombosis). The pathogenesis of persistent daily cephalgia is widely known, some studies believe that pain may be due to cytokine production and persistent glial activation, responsible for precipitating the clinical picture (50), also responsible for neurological disorders in COVID-19 (as we saw above).

Specific diseases, such as fibromyalgia, peripheral neuropathy or chronic migraine, have shown clinical modifications. Fibromyalgia has shown worse results in the evaluation tests (quality of sleep, fatigue, pain, rigidity) in patients suffering from COVID disease 19 (51). Later studies will confirm whether the worsening is due to the hypersensitivity of the disease. The study by Consonni and collaborators (52) observed during the pandemic that despite similar complaints about changing habits and concerns about the COVID-19 pandemic, patients with peripheral neuropathy and chronic migraine had different reactions. While, in patients with migraine, it affected behavior, mainly with psychological fragility. This suggests the need to personalize patient care based on their chronic pain conditions.

In conclusion, it takes time to confirm some of the data obtained in the few studies regarding the impact of COVID-19 disease

on chronic pain. Fundamentally confirming a physical basis, since the psychological aspect seems clearer, since even in people with a diagnosis of chronic pain not directly affected by the coronavirus, the pandemic outbreak and the subsequent blockade have been important stressors (53). Therefore, interdisciplinary interventions should be designed in which psychologists and health care staff work together to minimize the psychological distress of patients with chronic pain syndromes. These syndromes could increase the demand for medical care at an extremely difficult time. In this situation, telemedicine is an excellent option for patient management.

## REFERENCES

1. Wuhan Municipal Health Commission. Report of clustering pneumonia of unknown etiology in Wuhan City. 2019. <http://wjw.wuhan.gov.cn/front/web/showDetail/2019123108989>.
2. Declaración Universal de los Derechos Humanos. Available in: <http://www.un.org/es/documents/udhr> [Access 18 Feb 2021].
3. Pascarella G, Strumia A, Piliago C, Bruno F, Del Buono R Costa F, et al. COVID-19 diagnosis and management: a comprehensive review. *J Intern Med*. 2020;288(2):192-206. DOI: 10.1111/joim.13091.
4. Song XJ, Xiong DL, Wang ZY. Pain management during the COVID-19 pandemic in China: lessons learned. *Pain Med*. 2020;21(7):1319-23. DOI: 10.1093/pm/pnaa143.
5. Young BE, Ong SWX, Kalimuddin S. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *JAMA*. 2020;323:1488-94. DOI: 10.1001/jama.2020.3204.
6. Loeser JD, Treede RD. The Kyoto protocol of IASP Basic Pain Terminology. *Pain*. 2008;137:473-7. DOI: 10.1016/j.pain.2008.04.025.
7. Campbell A, Rodin R, Kropp R, Mao Y, Hong Z, Vachon J, et al. Risk of severe outcomes among patients admitted to hospital with pandemic (H1N1) influenza. *CMAJ*. 2010;182:349-55. DOI: 10.1503/cmaj.091823.
8. Hui DS, Chan MC, Wu AK, Ng PC. Severe acute respiratory syndrome (SARS): epidemiology and clinical features.

*The COVID-19 disease itself is associated with painful symptoms, including myalgia, arthralgia, abdominal pain, headache and chest pain*

- Postgrad Med J. 2004;80:373-81. DOI: 10.1136/pgmj.2004.020263.
9. Moldofsky H, Patcai J. Chronic widespread musculoskeletal pain, fatigue, depression and disordered sleep in chronic post-SARS syndrome; a case-controlled study. *BMC Neurol.* 2011;11:37. DOI: 10.1186/1471-2377-11-37.
  10. Hickie I, Davenport T, Wakefield D, Vollmer-Conna U, Cameron B, Vernon SD, et al. Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens: prospective cohort study. *BMJ.* 2006;333:575. DOI: 10.1136/bmj.38933.585764.AE.
  11. Garvin MR, Alvarez C, Miller JI, Prates ET, Walker AM, Amos BK, et al. A mechanistic model and therapeutic interventions for COVID-19 involving a RAS-mediated bradykinin storm. *Elife.* 2020;9:e59177. DOI: 10.7554/eLife.59177.
  12. Clauw DJ, Häuser W, Cohen SP, Fitzcharles MA. Considering the potential for an increase in chronic pain after the COVID-19 pandemic. *Pain.* 2020;161:1694-7. DOI: 10.1097/j.pain.0000000000001950.
  13. Beggs S, Liu XJ, Kwan C, Salter MW. Peripheral nerve injury and TRPV1-expressing primary afferent C-fibers cause opening of the blood-brain barrier. *Mol Pain.* 2010;6:74. DOI: 10.1186/1744-8069-6-74.
  14. Baig AM, Khaleeq A, Ali U, Syeda H. Evidence of the COVID-19 virus targeting the CNS: tissue distribution, host-virus interaction, and proposed neurotropic mechanisms. *ACS Chem Neurosci.* 2020;11:995-8. DOI: 10.1021/acschemneuro.0c00122.
  15. Tancheva L, Petralia MC, Miteva S, Dragomanova S, Solak A, Kalfin R, et al. Emerging neurological and psychobiological aspects of COVID-19 infection. *Brain Sci.* 2020;10(11):852. DOI: 10.3390/brainsci10110852.
  16. Gandhi S, Srivastava AK, Ray U, Tripathi PP. Is the collapse of the respiratory center in the brain responsible for respiratory breakdown in COVID-19 patients? *ACS Chem Neurosci.* 2020;11:1379-81.
  17. Guedj E, Million M, Dudouet P, Tissot-Dupont H, Bregeon F, Cammilleri S, et al. 18F-FDG brain PET hypometabolism in post-SARS-CoV-2 infection: substrate for persistent/delayed disorders? *Eur J Nucl Med Mol Imaging.* 2020;30:1-3.
  18. Marchand F, Perretti M, McMahon SB. Role of the immune system in chronic pain. *Nat Rev Neurosci.* 2005;6:521-32. DOI: 10.1038/nrn1700.
  19. Liu MM, Reidy AB, Saatee S. Perioperative steroid management: approaches based on current evidence. *Anesthesiology.* 2017;127:166-72. DOI: 10.1097/ALN.0000000000001659.
  20. Kosciuczuk U, Knapp P, Lotowska-Cwiklewska AM. Opioid-induced immunosuppression and carcinogenesis promotion theories create the newest trend in acute and chronic pain pharmacotherapy. *Clinics (Sao Paulo, Brazil).* 2020;75:1554. DOI: 10.6061/clinics/2020/e1554.
  21. Franchi S, Moschetti G, Amodeo G. Do all opioid drugs share the same immunomodulatory properties? A review from animal and human studies. *Front Immunol.* 2019;10:2914. DOI: 10.3389/fimmu.2019.02914.
  22. Saurer TB, Carrigan KA, Ijames SG. Suppression of natural killer cell activity by morphine is mediated by the nucleus accumbens shell. *J Neuroimmunol.* 2006;173:3-11. DOI: 10.1016/j.jneuroim.2005.11.009.
  23. McLean SA, Clauw DJ. Predicting chronic symptoms after an acute "stressor"-lessons learned from 3 medical conditions. *Med Hypotheses.* 2004;63:653-8. DOI: 10.1016/j.mehy.2004.03.022.
  24. Bendau A, Petzold MB, Pyrkosch L, Mascarell Maricic L, Betzler F, Rogoll J, et al. Associations between COVID-19 related media consumption and symptoms of anxiety, depression and COVID-19 related fear in the general population in Germany. *Eur Arch Psychiatry Clin Neurosci.* 2020 Jul 20;1-9. DOI: 10.1007/s00406-020-01171-6.
  25. Bakioğlu F, Korkmaz O, Ercan H. Fear of COVID-19 and Positivity: Mediating role of intolerance of uncertainty, depression, anxiety, and stress. *Int J Ment Health Addict.* 2020 May 28;1-14. DOI: 10.1007/s11469-020-00331-y.
  26. Chan D, Xu Feng Lin, Jane MG, Liu CW. Clinical challenges and considerations in management of chronic patients during a COVID-19 pandemic. *Ann Acad Med Singap.* 2020;49:669-73. DOI: 10.47102/annals-acadmedsg.2020130.
  27. Malhotra N, Joshi M, Datta R, Bajwa SJ, Mehdiratta L. Indian Society of Anaesthesiologists (ISA National) Advisory and

- Position Statement regarding COVID19. *Indian J Anaesth.* 2020;64:25963. DOI: 10.4103/ija.IJA\_288\_20.
28. Sarzi-Puttini P, Vellucci R, Zuccaro SM. The appropriate treatment of chronic pain. *Clin Drug Investig.* 2012;32:21-33. DOI: 10.2165/11630050-000000000-00000.
  29. Ghai B, Malhotra N, Jit S S. Telemedicine for chronic pain management during COVID-19 pandemic. *Indian J Anaesth.* 2020; 64:456-46. DOI: 10.4103/ija.IJA\_652\_20.
  30. Shanthanna H, Strand NH, Provenzano DA, Lobo CA, Eldabe S, Bhatia A, et al. Caring for patients with pain during the COVID-19 pandemic: Consensus recommendations from an international expert panel. *Anaesthesia.* 2020;75:935-44. DOI: 10.1111/anae.15076.
  31. Cohen SP, Baber ZB, Buvanendran A, McLean LTCBC, Chen Y, Hooten WM, et al. Pain management best practices from multispecialty organizations during the COVID-19 pandemic and public health crises. *Pain Med.* 2020;21:1331-46. DOI: 10.1093/pm/pnaa127.
  32. Chauhan V, Galwankar S, Arquilla B, Garg M, Somma SD, El-Menyar A, et al. Novel coronavirus (COVID-19): Leveraging telemedicine to optimize care while minimizing exposures and viral transmission. *J Emerg Trauma Shock.* 2020;13:20-4.
  33. Ayyagari A, Bhargava A, Agarwal R, Mishra SK, Mishra AK, Das SR, et al. Use of telemedicine in evading cholera outbreak in MahakumbhMela, Prayag, UP, India: An encouraging experience. *Telemed J E Health.* 2003;9:89-94. DOI: 10.1089/153056203763317693.
  34. Slattery BW, Haugh S, O'Connor L, Francis K, Dwyer CP, O'Higgins S, et al. An evaluation of the effectiveness of the modalities used to deliver electronic health interventions for chronic pain: Systematic review with network meta-analysis. *J Med Internet Res.* 2019;17:e1108. DOI: 10.2196/11086.
  35. Sundararaman LV, Edwards RR, Ross EL, Jamison RN. Integration of mobile health technology in the treatment of chronic pain: A critical review. *RegAnesth Pain Med.* 2017;42:488-98. DOI: 10.1097/AAP.0000000000000621.
  36. Fisher E, Law E, Dudeney J, Eccleston C, Palermo TM. Psychological therapies (remotely delivered) for the management of chronic and recurrent pain in children and adolescents. *Cochrane Database Syst Rev.* 2019;4:CD011118. DOI: 10.1002/14651858.CD011118.pub3.
  37. Cook AJ, Roberts DA, Henderson MD, Van Winkle LC, Chastain DC, Hamill-Ruth RJ. Electronic pain questionnaires: A randomized, crossover comparison with paper questionnaires for chronic pain assessment. *Pain.* 2004;110:310-7. DOI: 10.1016/j.pain.2004.04.012.
  38. Azevedo BR, Oliveira CB, Araujo GMD, Silva FG, Damato TM, Pinto RZ, et al. Is there equivalence between the electronic and paper version of the questionnaires for assessment of patients with chronic low back pain? *Spine.* 2020;45:E329-35. DOI: 10.1097/BRS.0000000000003281.
  39. Chhabra HS, Sharma S, Verma S. Smartphone app in self-management of chronic low back pain: A randomized controlled trial. *Eur Spine J.* 2018;27:2862-74. DOI: 10.1007/s00586-018-5788-5.
  40. Flodgren G, Rachas A, Farmer AJ, Inzitari M, Shepperd S. Interactive telemedicine: Effects on professional practice and health care outcomes. *Cochrane Database Syst Rev.* 2015;2015:CD002098. DOI: 10.1002/14651858.CD002098.pub2.
  41. Baker J, Stanley A. Telemedicine technology: A review of services, equipment, and other aspects. *Curr Allergy Asthma Rep.* 2018;18:60. DOI: 10.1007/s11882-018-0814-6.
  42. Boodley CA. Primary care telehealth practice. *J Am Acad Nurse Pract.* 2006;18:343-5. DOI: 10.1111/j.1745-7599.2006.00146.x.
  43. Aguarwal N, Jain P, Pathak R, Gupta R. Telemedicine in India: A tool for transforming health care in the era of COVID-19 pandemic. *J Educ Health Promot.* 2020;9:190. DOI: 10.4103/jehp.jehp\_472\_20.
  44. Lovell N, Maddocks M, Etkind SN, Taylor K, Carey I, Vora V, et al. Characteristics, symptom management, and outcomes of 101 patients with COVID-19 referred for hospital palliative care. *J Pain Symptom Manage.* 2020;60:77-81. DOI: 10.1016/j.jpainsymman.2020.04.015.
  45. Kemp HI, Corner E, Colvin LA. Chronic pain after COVID-19: implications for rehabilitation. *Br J Anaesth.* 2020;125:436-40. DOI: 10.1016/j.bja.2020.05.021.
  46. Niazkar HR, Zibae B, Nasimi A, Bahri N. The neurological manifestations of COVID-19: a review article. *Neurol Sci.* 2020;41:1667-71. DOI: 10.1007/s10072-020-04486-3.



47. Nieto R, Pardo R, Sora B, Feliu-Soler A, Luciano JV. Impact of COVID-19 lockdown measures on spanish people with chronic pain: an online study survey. *J Clin Med*. 2020;9:3558. DOI: 10.3390/jcm9113558.
48. Uygun Ö, Ertaş M, Ekizoğlu E, Bolay H, Özge A, Orhan EK, et al. Headache characteristics in COVID-19 pandemic-a survey study. *J Headache Pain*. 2020;21:121. DOI: 10.1186/s10194-020-01188-1.
49. Sampaio Rocha-Filho PA, Voss L. Persistent headache and persistent anosmia associated with COVID-19. *Headache*. 2020;60:1797-9. DOI: 10.1111/head.13941.
50. Yamani N, Olesen J. New daily persistent headache: a systematic review on an enigmatic disorder. *J Headache Pain*. 2019;20:80. DOI: 10.1186/s10194-019-1022-z.
51. Salaffi F, Giorgi V, Sirotti S, Bongiovanni S, Farah S, Bazzichi L, et al. The effect of novel coronavirus disease-2019 (COVID-19) on fibromyalgia syndrome. *Clin Exp Rheumatol*. 2020;16.
52. Consonni M, Telesca A, Grazzi L, Cazzato D, Lauria G. Life with chronic pain during COVID-19 lockdown: the case of patients with small fibre neuropathy and chronic migraine *Neurol Sci*. 2021;42(2):389-97.
53. Serrano-Ibáñez ER, Esteve R, Ramírez-Maestre C, Ruiz-Párraga GT, López-Martínez AE. Chronic pain in the time of COVID-19: Stress aftermath and central sensitization. *Br J Health Psychol*. 2020. DOI: 10.1111/bjhp.12483. Online ahead of print